Author index

Abdel-Rahim, A.M. *Phoma sorghina* causing a leaf spot of water hyacinth in the Sudan, 429

Adams, S.E., Jones, R.A.C. and Coutts, R.H.A. Occurrence of resistance-breaking strains of potato virus X in potato stocks in England and Wales, 435

Adena, M.A., see Guy *et al.*, 337

Allen, Elizabeth C., see Jordan, 547

Archer, S.A., see Woolacott, 225

Asher, M.J.C. and Thomas, C.E. Components of partial resistance to *Erysiphe graminis* in spring barley, 123

Ayres, P.G., see Paul, N.D., 561

Bailliss, K.W. and Senananyaka, S. Virus infection and reproductive losses in faba bean (*Vicia faba* L.), 185

Bandara, J.M.R.S., see Randombage, 479

Ball, S.L., see Idris, 219

Baulcombe, D., Flavell, R.B., Boulton, R.E. and Jellis, G.J. The sensitivity and specificity of a rapid nucleic acid hybridization method for the detection of potato virus X in crude sap samples, 361

Bennett, Fiona G.A. Resistance to powdery mildew in wheat: a review of its use in agriculture and breeding programmes, 279

Bevercombe, G.P. and Rayner, A.D.M. Population structure of *Cryptostroma corticale*, the causal fungus of sooty bark disease of sycamore, 211

Black, R., see Lumyong, 431

Blakeman, J.P., see Parker, 71

Boulton, R.E., see Baulcombe *et al.*, 361

Bradshaw, N.J. *Septoria cucurbitacearum* on courgettes – a new British record, 135

Brown, M.C., Taylor, G.S. and Epton, H.A.S. Carbendazim resistance in the eyespot pathogen *Pseudocercosporella herpotrichoides*, 101

Burrell, M.M. Inhibition of browning, phenoxyacetic acids and phenolic metabolism in potato tuber discs: a model system to study chemicals that control common scab, 325

Channon, A.G., see Holmes, 347

Chase, A.R. *Xanthomonas campestris* pv. *hederae* causes a leaf spot of five species of Araliaceae, 439

Chen, W.Y. and Echandi, E. Effects of avirulent bacteriocin-producing strains of *Pseudomonas solanacearum* on the control of bacterial wilt of tobacco, 245

Chin, K.M. and Wolfe, M.S. The spread of *Erysiphe graminis* f. sp. *hordei* in mixtures of barley varieties, 89

Channon, A.G. and Hissett, R. The incidence of bacterial wilt caused by *Xanthomonas campestris* pv *graminis* in pasture grasses in the West of Scotland, 113

Chin, K.M., Wolfe, M.S. and Minchin, P.N. Host-mediated interactions between pathogen genotypes, 161

Chin, K.M. and Wolfe, M.S. Selection on *Erysiphe graminis* in pure and mixed stands of barley, 535

Christian, D.G. and Miller, D.P. *Cephalosporium* stripe in winter wheat grown after different methods of straw disposal, 605

Cockbain, A.J., see Jones, 599

Cole, J.S., see Winch *et al.*, 467
Coley-Smith, J.R., see Esler, 199
Coley-Smith, J.R., see Parfitt *et al.*, 441
Cooper, J.I., see Massalski, 255
Cooper, R.M., see Street, 483
Couch, H.B., see Muchovej, 589
Coutts, R.H.A., see Adams *et al.*, 435
Crompton, J.G., see Kirk, 317
Crosse, J.E., see Kairu *et al.*, 131
Cross, J.V. Biological control of two-spotted spider mite (*Tetranychus urticae*) by *Phytoseiulus persimilis* on strawberries grown in 'walk-in' plastic tunnels, and a simplified method of spider mite population assessment, 417
Crute, I.R., see Michelmore *et al.*, 301
Crute, I.R., see Horwood, 385

Dale, J.L., see Guy *et al.*, 337
Deverall, B.J., see Sutton, 337
Dickinson, C.H., see Hims *et al.*, 513

Echandi, E., see Chen, 245
El-Khadam, M. and Papavizas, G.C. Effect of the herbicides EPTC and linuron on cotton diseases caused by *Rhizoctonia solani* and *Fusarium oxysporum* f. sp. *vasinfectum*, 411
Epton, H.A.S., see Brown *et al.*, 101
Esler, G. and Coley-Smith, J.R. Resistance to *Sclerotium cepivorum* in Allium and other genera, 199

Flavell, R.B., see Baulcombe *et al.*, 361
Fletcher, J.T., see Hims *et al.*, 513
Forrest, J.M.S. and Phillips, M.S. The effect of continuous rearing of a population of *Globodera pallida* (Pa 2) on susceptible or partially resistant potatoes, 53
Freigoun, S.O., see Jones *et al.*, 601
French, N., see Port, 133
Frost, R.R., see Hicks, 581
Fuller, Nicola J., see Jones, 595

Gibbs, A.J., see Guy *et al.*, 337
Gibson, R.W., see Katis, 425
Guile, C.T. Damage to rhubarb (*Rheum rhaponticum* L.) and Virginia creeper (*Parthenocissus quinquefolia* (L.) Planch.) by dock beetle (*Gastrophysa viridula* Degeer), 269
Guy, P.L., Dale, J.L., Adena, M.A. and Gibbs, A.J. A taxonomic study of the host range of tymoviruses, 337

Hall, K. and Kavanagh, J.A. Laboratory studies on the growth and reproduction of *Cladosporium allii-cepae*, the cause of leaf blotch of onion, 147
Heale, J.B., see Wright, 493
Hicks, R.G.T. and Frost, R.R. Isolation of a tobamovirus from rose, 581
Hill, S.A. and Jackson, Elizabeth A. An investigation of the reliability of ELISA as a practical test for detecting potato leaf roll virus and potato virus Y in tubers, 21

Hill, S.A., see Wright *et al.*, 591

Hims, M.J., Dickinson, C.H. and Fletcher, J.T. Control of red thread, a disease of grasses caused by *Laetisaria fuciformis*, 513

Hissett, R., see Channon, 113

Holden, D.W. and Strange, R.N. Biochemical characteristics of sporelings of *Puccinia graminis* f. sp. *tritici* and protoplasts from wheat isolines of potential significance in specificity, 233

Hollis (Jr), J.P. The Horsfall-Barratt Grading System, 145

Holmes, S.J.I. and Channon, A.G. Studies on metalaxyl-resistant *Phytophthora infestans* in potato crops in south-west Scotland, 347

Hollomon, D.W. A laboratory assay to determine the sensitivity of *Rhynchosporium secalis* to the fungicide triadimenol, 65

Humpherson-Jones, F.M., see Maude *et al.*, 525

Idris, M.O. and Ball, S.L. Inter and intracontinental sexual compatibility in *Sclerospora graminicola*, 219

Ingram, D.S., see Michelmore, 301

Inwang, Bunjerd, see Lumyong *et al.*, 431

Jackson, Elizabeth A., see Hill, 21

Jackson, A.W., see Wardlow, 57

Jackson, G.V.H., see Winch *et al.*, 467

Javed, Z.U.R. Control of benomyl-resistant strains of *Colletotrichum coffeanum*, the cause of coffee berry disease, 27

Javed, Z.U.R., see Kairu *et al.*, 131

Jeger, M.J. Relating disease progress to cumulative numbers of trapped spores: apple powdery mildew and scab epidemics in sprayed and unsprayed orchard plots, 517

Jellis, G.J., see Baulcombe *et al.*, 361

Jones, R.A.C., see Adams *et al.*, 435

Jones, R.A.C. and Fuller, Nicola J. Incidence of potato virus V in potato stocks in England and Wales, 595

Jones, P., Cockbain, A.J. and Freigoun, S.O. Association of a mycoplasma-like organism with broad bean phyllody in the Sudan, 599

Jordan, V.W.L. and Allen, Elizabeth C. Barley net blotch: influence of straw disposal and cultivation methods in inoculum potential, and on incidence and severity of autumn disease, 547

Kairu, G.M., Nyangena, C.M.S., Javed, Z.U.R. and Crosse, J.E. Iatrogenic effects of captafol on bacterial blight of coffee, 131

Katis, N. and Gibson, R.W. Transmission of beet mosaic virus by cereal aphids, 425

Kavanagh, J.A., see Hall, 147

Kirk, P.M. and Crompton, J.G. Pathology and taxonomy of *Cladosporium* leaf blotch of onion (*Allium cepa*) and leek (*A. porrum*), 317

Lapwood, D.H., Read, P.J. and Spokes, Janis. Methods for assessing the susceptibility of potato tubers of different cultivars to rotting by *Erwinia carotovora* subspecies *atroseptica* and *carotovora*, 13

Le Cuirot, Carolyne, see Wright, 591

Lumyong, Pipob, Black, R. and Inwang, Bunjerd. *Embellisia abundans* causing leaf spot of highland-grown Brussels sprouts in Northern Thailand, 431

Lyshol, A.J., Semb, L. and Taksdal, G. Reduction of cavity spot and root dieback in carrots by fungicide applications, 193

MacNish, G.C. The use of undisturbed soil cores to study methods of controlling rhizoctonia patch of cereals, 355

McCracken, A.R. *Pythium paroecandrum* associated with a root-rot of parsley, 603

McGregor, A.J. Comparison of cuprous oxide and metalaxyl with mixtures of these fungicides for the control of *Phytophthora* pod rot of cocoa, 81

McHale, T., see Parfitt *et al.*, 441

Mansfield, J.W., see Stewart, 401

Massalski, P.R. and Cooper, J.I. The location of virus-like particles in the male gametophyte of birch, walnut and cherry naturally infected with cherry leaf roll virus and its relevance to vertical transmission of the virus, 255

Maude, R.B., Shipway, M.R., Presly, A.H. and O'Connor, D. The effects of direct harvesting and drying systems on the incidence and control of neck rot (*Botrytis allii*) in onions, 263

Maude, R.B., Humpherson-Jones, F.M. and Shuring, Catriona, G. Treatments to control *Phoma* and *Alternaria* infections of brassica seeds, 525

Michelmore, R.W., Norwood, J.M., Ingram, D.S., Crute, I.R. and Nicolson, P. The inheritance of virulence in *Bremia lactucae* to match resistance factors 3, 4, 5, 6, 8, 9, 10 and 11 in lettuce (*Lactuca sativa*), 301

Miller, D.P., see Christian, 607

Minchin, P.N., see Chin, 161

Muchovej, J.J. and Couch, H.B. *Marasmius graminum*, a colonizer of senescent leaves of Kentucky bluegrass (*Poa pratensis*), 589

Newhook, F.J., see Winch *et al.*, 467

Newman, Pamela L. Differential host-parasite interactions between oilseed rape and *Leptosphaeria maculans*, the causal fungus of stem canker, 205

Nicholson, P., see Michelmore, 301

Norwood, J.M., see Michelmore, 301

Norwood, Judith M. and Crute, I.R. The genetic control and expression of specificity in *Bremia lactucae* (lettuce downy mildew), 385

Nyangena, C.M.S., see Kairu, 131

O'Connor, D., see Maude *et al.*, 263

Papavizas, G.C., see El-Khadem, 411

Parfitt, D., Coley-Smith, J.R. and McHale, T. *Laterispora brevirama* on sclerotia of *Sclerotinia*, 441

Parker, Ann and Blakeman, J.P. Nutritional factors to determine the sensitivity of *Rhynchosporium secalis* to the fungicide triadimenol, 71

Paul, N.D. and Ayres, P.G. Effects of rust and post-infection drought on photosynthesis, growth and water relations in groundsel, 561

Pearson, M.N. and Robb, Sheila, M. The occurrence and effects of barley yellow dwarf virus in maize in SW England, 503

Phillips, M.S., see Forrest, 53

Port, C.M. and French, N. Damage to spring barley by larvae of *Dilophus febrilis* (L.) (Bibionidae: Diptera), 133

Presly, A.H., see Maude, 263

Presly, A.H. Retention of iprodione on salad onion leaves in relation to the control of *Botrytis* spp., 571

Randombage, Sunietha and Bandara, J.M.R.S. Little leaf disease of *Piper nigrum* in Sri Lanka, 479

Rayner, A.D.M., see Bevercombe, 211

Read, P.J., see Lapwood, 13

Robb, Sheila M., see Pearson, 503

Semb, L., see Lyshol, 193

Senananyake, S., see Bailiss, 185

Sewell, G.W.F. and Wilson, J.F. The nature and distribution of *Verticillium albo-atrum* strains highly pathogenic to the hop, 39

Sharma, H.S.S. Assessment of the reaction of some spring barley cultivars to *Pyrenophora teres* using whole plants, detached leaves and toxin bioassay, 371

Shattock, R.C., see Skidmore *et al.*, 173

Shaw, D.S., see Skidmore *et al.*, 173

Shipway, M.R., see Maude *et al.*, 263

Shuring, Catriona G., see Maude *et al.*, 525

Skidmore, D.I., Shattock, R.C. and Shaw, D.S. Oospores in cultures of *Phytophthora infestans* resulting from selfing induced by the presence of *P. drechsleri* isolated from blighted potato foliage, 173

Spokes, Janis, see Lapwood *et al.*, 13

Stewart, Alison and Mansfield, J.W. Fungal development and plant response in detached onion bulb scales and leaves inoculated with *Botrytis allii, B. cinerea, B. fabae* and *B. squamosa*, 401

Strange, R.N., see Holden, 233

Street, P.F.S. and Cooper, R.M. Quantitative measurement of vascular flow in petioles of healthy and Verticillium-infected tomato, 483

Sutton, D.C. and Deverall, B.J. Phytoalexin accumulation during infection of bean and soybean by ascospores and mycelium of *Sclerotinia sclerotiorum*, 377

Taksdal, G., see Lyshol, 193

Taylor, G.S., see Brown, 101

Thomas, C.E., see Asher, 123

Thomas, B.J. Rose mosaic disease: symptoms induced in roses by graft inoculation with both prunus necrotic ringspot and apple mosaic viruses, 155

Wardlow, L.R. and Jackson, A.W. Comparison of laboratory methods for assessing numbers of apple rust mite (*Aculus schlechtendali*) overwintering on apple, 57

Wastie, R.L. Inoculating plant material by jet injection, 61

Whitehead, A.G. Interaction of three lucerne cultivars and eleven English isolates of stem nematode (*Ditylenchus dipsaci*), 'lucerne race', 33

Wilson, J.F., see Sewell, 39

Winch, J.E., Newhook, F.J., Jackson, G.V.H. and Cole, J.S. Studies of *Colletotrichum gloeosporioides* disease on yam, *Dioscorea alata*, in Solomon Islands, 467

Wolfe, M.S., see Chin, 89

Wolfe, M.S., see Chin *et al.*, 161

Wolfe, M.S. Trying to understand and control powdery mildew, 451

Wolfe, M.S., see Chin, 535

Wood, R.K.S. Establishment of infection, 3

Woolacott, B. and Archer, S.A. The influence of the primary gem tube on infection of barley by *Erysiphe graminis* f. sp. *hordei*, 225

Wright, Alison, J. and Heale, J.B. Adult plant resistance to powdery mildew (*Erysiphe graminis*) in three barley cultivars, 493

Wright, Daphne M., Le Cuirot, Carolyne and Hill, S.A. Identification of Zucchini Yellow Mosaic Virus from courgettes in Jersey by immune electron microscopy, 591